



## CLINICAL STUDY

# EVALUATION OF POSTURAL STABILITY AND FALL RISK IN PATIENTS WITH OBSTRUCTIVE SLEEP APNEA SYNDROME

Harun SOYALIÇ<sup>1</sup>, MD; Handan İNÖNÜ<sup>2</sup>, MD; Mustafa AVCU<sup>1</sup>, MD; Can KOÇAK<sup>3</sup>, MD; Ahmet İNANIR<sup>4</sup>, MD; Evrim Ünsal TUNA<sup>1</sup>, MD

<sup>1</sup>Ahi Evran Üniversitesi Tıp Fakültesi, Kulak Burun Boğaz Anabilim Dalı, Kırşehir, Turkey <sup>2</sup>Gaziosmanpaşa Üniversitesi Tıp Fakültesi, Göğüs Hastalıkları Anabilim Dalı, Tokat, Turkey <sup>3</sup>Gaziosmanpaşa Üniversitesi Tıp Fakültesi, Kulak Burun Boğaz Anabilim Dalı, Tokat, Turkey <sup>4</sup>Gaziosmanpaşa Üniversitesi Tıp Fakültesi, Fizik Tedavi ve Rehabilitasyon Anabilim Dalı, Tokat, Turkey

### SUMMARY

**Objectives:** To evaluate patients with obstructive sleep apnea syndrome (OSAS) for fall risk and postural stability using the Biodex Balance System.

**Methods:** The study involved the examination of 95 individuals. Twenty-five of these individuals were healthy controls (13 male, 12 female) whose apnea hypopnea index (AHI) was below 5. In addition, 35 of them were patients with mild OSAS (17 male, 18 female); and 35 of them were patients with moderate-severe OSAS (19 male, 16 female). OSAS patients and healthy controls were examined for postural stability and fall risk using the Biodex Balance System.

**Results:** No statistically significant difference was observed between groups regarding sex, age, height, weight, Overall Stability Index, Antero-Posterior Stability Index, or Medio-Lateral Stability Index ( $p>0.05$ ). The mean body mass index (BMI) in the control group did not show a significant difference in comparison to the mild OSAS or moderate-severe OSAS groups ( $p>0.05$ ). Interestingly, the mean BMI of the mild OSAS group ( $27.11\pm 2.35$  kg/m<sup>2</sup>) was lower compared to the moderate-severe OSAS group ( $28.72\pm 2.98$  kg/m<sup>2</sup>) ( $p=0.039$ ). Furthermore, a significant difference was observed between the groups for Fall Risk Index (FRI) ( $p<0.001$ ). The moderate-severe OSAS group had a significantly higher FRI compared to both the mild OSAS group and the control group ( $p=0.044$ ,  $p<0.001$ , respectively).

**Conclusion:** Postural instability and fall risk in patients with severe OSAS that engage in activities that require attention-balance can be assessed using the Biodex Balance System.

**Keywords:** Biodex balance system, postural instability, fall risk, obstructive sleep apnea syndrome

### OBSTRÜKTİF UYKU APNESİ OLAN HASTALARDA POSTURAL DENGE VE DÜŞME RİSKİNİN DEĞERLENDİRİLMESİ

#### ÖZET

**Giriş:** Biodeks Denge Sistemini kullanarak obstrüktif uyku apneli hastalarda postural dengenin ve düşme riskinin (Postural Stability and Fall Risk) değerlendirilmesidir.

**Yöntem ve Gereçler:** Çalışmaya 95 birey dahil edildi. Bireylerin 25'i AHI'si 5 in altında olan sağlıklı kontrol grubundan (13 erkek, 12 kadın), 35'i hafif dereceli OUAS 'lı hastalardan (17 erkek, 18 kadın), 35'i de orta -ağır OUAS'lı hastalardan (19 erkek, 16 kadın) oluşmaktaydı. OUAS 'lı hastalar ve sağlıklı kontrol grubu bireyler denge ve düşme riski açısından test edildi. Denge ve düşme riski Biodeks Denge Sistemi ile değerlendirildi.

**Bulgular:** Grupların cinsiyet, yaş, boy, kilo, Genel Stabilité İndeksi (GSİ), Antero-Posterior Stabilité İndeksi (APSi) ve Medio-Lateral Stabilité İndeksi (MLSİ) arasında karşılaştırma yapıldığında istatistiksel olarak anlamlı bir farklılık saptanmadı ( $p>0.05$ ). Kontrol grubu ile hafif OUAS ve orta- ağır OUAS gruplarının VKİ ortalamaları arasında anlamlı bir fark saptanmaz iken ( $p>0.05$ ) hafif OUAS grubunun VKİ ortalaması ( $27,11\pm 2,35$ ) orta-ağır ağır OUAS grubunun VKİ ortalamasından ( $28,72\pm 2,98$ ) anlamlı derecede daha düşüktü ( $p=0,039$ ). Gruplar Düşme Riski İndeksine (DRİ) göre karşılaştırıldığında gruplar arasında istatistiksel olarak anlamlı bir farklılık saptandı ( $p<0,001$ ). Orta-ağır OUAS grubunun ortalama Düşme Riski İndeksi, hafif OUAS grubunun ortalama Düşme Riski İndeksi ve kontrol grubunun ortalama Düşme Riski İndeksi değerinden istatistiksel olarak anlamlı derecede daha büyüktü. (sırayla  $p=0,044$ ;  $p<0,001$ ).

**Sonuç:** Özellikle ağır derecede obstrüktif uyku apnesi olan ve dikkat-denge gerektiren aktiviteler gerçekleştiren hastalardaki dengesizlik ve düşme riski Biodeks Denge Sistemi ile belirlenebilir.

**Anahtar Sözcükler:** Biodeks denge sistemi, denge bozukluğu, düşme riski, obstrüktif uyku apnesi sendromu

## INTRODUCTION

Obstructive sleep apnea syndrome (OSAS), a sleep disorder that affects adults, has been described by sleep-associated collapse of the upper airway.

Corresponding Author: Harun Soyaliç MD Ahi Evran Üniversitesi Tıp Fakültesi, Kulak Burun Boğaz Anabilim Dalı, Kırşehir, Turkey, E-mail: harun\_soyalic@hotmail.com

Received: 10 July 2017, revised for: 03 August 2017, accepted for publication: 07 August 2017

OSAS has been linked to several symptoms such as snoring, fatigue, concentration or memory lapses, and excessive daytime sleepiness, and cardiopulmonary dysfunctions<sup>1</sup>.

OSAS has been demonstrated to impact free radical and oxidative stress factor levels as well as ischemic conditions, all of which alter normal central and peripheral nervous system function. Oxygen



desaturation/resaturation in OSAS has been shown to elicit inflammatory components, oxidative damage, and endothelium dysregulation. Multiple reports have shown that vasa nervorum damage caused by insufficient oxygenation may result in damage peripheral nerves and the central nervous system<sup>2,3</sup>.

Recent studies have reported an association between OSAS and idiopathic dizziness<sup>4,5,6,7,8</sup>. Interestingly, vestibular nuclei have been shown to be more sensitive to hypoxia than other cerebral nuclei<sup>9</sup>. Furthermore, the functional consequences associated with vestibular nuclei changes may be an important index of abnormal respiratory nuclei activity<sup>8</sup>.

Postural balance control is a complex motor ability that requires flexibility in addition to organization of sensory inputs. Environmental components influencing balance include vestibular, somatosensorial and visual systems, and inputs, all of which are united in the central nervous system to determine numerous appropriate muscular responses for controlling body position and posture on a supporting base. Maintenance of postural stability requires an intact neuromuscular system, an effective motor response, and adequate supporting muscle power to restore balance when it is disturbed<sup>10,11,12</sup>.

Weak muscles, vertigo-dizziness, visual problems, drug side effects, and environmental factors are the main causes for falls. Patients with OSAS, particularly with a moderate-severe degree, have significant risks in their daily life that require attention. For assessment of fall risk, dynamic tests are more efficient than static tests. Therefore, it has been advised that dynamic tests (Biodex Balance System) that are appropriate for locomotor tasks should also be performed during assessment of postural stability<sup>13,14,15</sup>.

As a laboratory assessment method for postural stability, the Biodex Balance System (BBS) has been described for its reliability in assessing dynamic postural stability in healthy and visually-impaired individuals (Intraclass Correlation Coefficient (ICC) ranges from 0.59 to 0.95)<sup>16,17</sup>.

The goal of this report was to evaluate the fall risk and postural stability OSAS patients diagnosed clinically and with polysomnography using the BBS.

## MATERIAL and METHODS

### 1. Subjects

The study included 95 individuals. Of these patients, 46 (48.4%) were female, and 49 (51.6%) were male. Twenty-five (13 males, 12 females) individuals with an apnea hypopnea index (AHI)

below 5 constituted the healthy control group; 35 (17 males, 18 females) patients constituted mild OSAS group; and 35 (19 males, 16 females) patients constituted moderate-severe OSAS group.

Informed consent is in accordance with the study protocol, approved by the ethics committee of Gaziosmanpaşa Medical Faculty (12-KAEK-126)

The study exclusion criteria were the presence of a disease that may cause central or peripheral vertigo such as chronic neurological disease, Meniere's disease, perilymph fistula, chronic middle ear infection and migraine, alcohol or substance abuse or use of any drugs that can impair balance, and presence of a chronic disease (rheumatic diseases, muscular diseases, malignancy, hepatitis, and hypertension).

### 2. Study design

Fall risk and postural stability was assessed using the Biodex Balance System [(BBS) Biodex Inc., New York]. The BBS comprises of a mobile balance platform controlled by a computer software that allows up to 20° tilt of the surface, which enables objective evaluation of balance. Overall Stability Index (OSI), Antero-Posterior Stability Index (APSI), Medio-Lateral Stability Index (MLSI), and Fall Risk Index (FRI) can be assessed by using this system. High scores from these tests mean impaired balance and increased fall risk. The tests were performed by identifying the coordination of subjects were standing on two bare feet on the platform, while knees were slightly flexed (10-15°), taking the most appropriate position for maintaining balance, and when eyes were open. All participants were informed about the tests and the rules that he/she must obey during the test. For postural stability and fall risk, three tests were performed, and each lasted for about 20 seconds. The participants' weight, height, GSI, APSI, MLSI, and FRI data were evaluated.

### 3. Statistical analysis

Data analysis was performed using SPSS 22.0 (IBM Corporation, New York, United States) software. Normality assessment was made using Shapiro-Wilk test, and a Levene test was used to test the variance homogeneity. Comparison of more than two groups for numerical data was made using a one-way ANOVA test for parametric data, and a Sidak test was used for post hoc analysis. For non-parametric data, the Kruskal Wallis H Test Monte Carlo simulation analysis was performed, while a Dun's test was used for post hoc analysis.



Statistical analyses were performed within 95% confidence level, and a p-value of less than 0.05 was accepted as significant.

## RESULTS

The mean patient age was 43.71±9.00 years, the maximum age was 62 years, and the minimum age was 21 years.

The mean body mass index (BMI) of the control group was not significantly different from the mild OSAS group and severe OSAS group (p>0.05). However, the mean BMI of the mild OSAS group (27.11±2.35 kg/m<sup>2</sup>) was significantly lower in comparison to the severe OSAS group (28.72±2.98 kg/m<sup>2</sup>) (p=0.039).

There was no statistically significant difference between the groups regarding distribution of sex, age, height, weight, Overall Stability Index (OSI), Antero-Posterior Stability Index (APSI), and Medio-Lateral Stability Index (MLSI) (p>0.05) (Table 1).

There was a statistically significant difference between the groups when compared for Fall Risk Index (FRI) (p<0.001) (Figure 1). The mean FRI of moderate-severe OSAS group was significantly higher compared to both the mild OSAS group and the control group (p=0.044 and p<0.001, respectively).

**Table 1:** The evaluation of control, mild OSAS, and severe OSAS groups according to sex, age, height, weight, BMI, AHI, general stability index, anterior posterior stability index, medio-lateral stability index, and fall risk. Regarding comparison of the groups in terms of Fall Risk Index (FRI), there was a statistically significant difference between the groups (p<0.001). The mean FRI in the moderate-severe OSAS group was significantly higher compared to the mild OSAS group and the control group (p=0.044 and p<0.001, respectively).

	Control=I	Mild OSAS=II	Moderate-Severe OSAS=III	Total	P Value	Pairwise Comparison		
	(n=25)	(n=35)	(n=35)	(n=95)		I-II	I-III	II-III
<b>Gender</b>	<b>n(%)</b>	<b>n(%)</b>	<b>n(%)</b>	<b>n(%)</b>				
Female	12 (48.0)	18 (51.4)	16 (45.7)	46 (48.4)	0.965	ns	ns	ns
Male	13 (52.0)	17 (48.6)	19 (54.3)	49 (51.6)				
	<b>Mean±SD.</b>	<b>Mean±SD.</b>	<b>Mean±SD.</b>	<b>Mean±SD.</b>				
Age	45.20±8.08	44.66±8.24	41.69±10.15	43.71±9.00	0.243	ns	ns	ns
Height	166.88±6.55	167.26±6.76	167.00±6.19	167.06±6.43	0.973	ns	ns	ns
Weight	76.76±9.85	75.97±9.12	80.14±9.88	77.72±9.68	0.168	ns	ns	ns
Body Mass Index	27.50±2.60	27.11±2.35	28.72±2.98	27.80±2.73	<b>0.037</b>	0.922	0.233	<b>0.039</b>
	<b>Med. (Max. / Min.)</b>	<b>Med. (Max. / Min.)</b>	<b>Med. (Max. / Min.)</b>	<b>Med. (Max. / Min.)</b>				
Apnea hipopnea Index	2 (8 / 0.2)	9.5 (15 / 5.1)	36 (105 / 15)	11.6 (105 / 0.2)	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>
General Stability Index	0.6 (1.2 / 0.2)	0.5 (3.2 / 0.3)	0.5 (3.2 / 0.2)	0.5 (3.2 / 0.2)	0.873	ns	ns	ns
Antero-Posterior Stability Index	0.4 (0.8 / 0.2)	0.4 (2.7 / 0.2)	0.4 (2.9 / 0.1)	0.4 (2.9 / 0.1)	0.709	ns	ns	ns
Medio-Lateral Stability Index	0.3 (0.8 / 0.1)	0.2 (1.6 / 0.1)	0.2 (0.8 / 0.1)	0.2 (1.6 / 0.1)	0.277	ns	ns	ns
Fall Risk Index	1.5 (2.4 / 0.4)	1.8 (5 / 0.4)	2.3 (6.6 / 0.9)	2 (6.6 / 0.4)	<b>&lt;0.001</b>	0.217	<b>&lt;0.001</b>	<b>0.044</b>

Pearson Chi Square Test (Monte Carlo) / Oneway ANOVA Test Post Hoc Test: Sidak / Kruskal Wallis Test (Monte Carlo) Post Hoc Test : Dunn's Test / SD.:Standard Deviation - Med.:Median - Max.:Maximum - Min.:Minimum - ns:Not Significant




Figure 1: Comparison of control, mild OSAS, and moderate-severe OSAS groups based on their fall risk.

## DISCUSSION

In patients with OSAS, hypoxia-induced impairment of peripheral and central vestibular has been documented for its association with dizziness and vertigo. The cardiovascular and neurological complications caused by hypoxia in OSAS has been well characterized. However, the consequences that arise from neurological damage have not been clearly defined<sup>8</sup>.

The functional consequences associated with vestibular nuclei changes may be an important index of abnormal respiratory nuclei activity. Interestingly, vestibular nuclei damage has been revealed previously by observing any changes in the slow and rapid eye movements<sup>8</sup>.

Experimental OSAS studies in rats have shown that severe OSAS may affect the nervous system by inducing neural loss in the nucleus ambiguus<sup>4</sup>.

Periodic ischemia/reperfusion that is associated with OSAS results causes many complications including inflammation, endothelial dysfunction, and oxidative stress. These complications can further affect the sympathetic and parasympathetic nervous systems, hematopoiesis, and coagulation. Early in ischemia damage, mechanisms to protect peripheral neuropathy are triggered, but become inadequate and neuropathy prevails in chronic hypoxemia<sup>18,19</sup>.

It is reported that the postural instability causes the impaired balance and this condition is also

related to the increased risk of falling. Although there are many factors affecting fall risk such as walking problems, loss of muscular activity, and drugs, there is a strong relation between postural instability and increased risk of falls. Falling phobia restricts many activities. This condition can lead to a decrease in quality of life<sup>20,21,22</sup>.

The BBS was introduced in 1999, and is a reliable method for the assessment of fall risk and postural stability in various diseases. Several studies have used BBS to evaluate fall risk and postural stability in ankylosing spondylitis and rheumatoid arthritis patients<sup>20,21,22</sup>. These patients have been shown to have increased fall risk. In our study, Although postural stability indexes of moderate-severe OSAS group, OSI, APSI, and MLSI were higher than the control group, we did not detect any statistically significant impairment in the postural stability tests of patients with OSAS. However, we found that The FRI was statistically significant. increased in patients with moderate-severe OSAS. It can be said that it indicates there is a predisposition to a impairment balance.

Multiple reports have evaluated the relationship between vertigo-dizziness and OSAS with contradicting results<sup>4,5,8,23,24,25</sup>. It was shown that sleep deprivation in humans can cause disturbances in the posterior parietal cortex, which is known for its role in the processing of vestibular information and mediation of vestibulo-ocular reflex (VOR)<sup>25</sup>.



Vestibular dysfunction has been shown to be more pronounced in moderate-severe OSAS patients in comparison to patients with mild OSAS<sup>8,23</sup>.

Sensitivity of the posterior labyrinth to a hypoxic state was proven with caloric stimulation tests and showed changes in the vestibular reflex. Patients with severe OSAS had worse results in this test. Considering the sensitivity of the posterior labyrinth to a hypoxic state and the anatomical proximity of these two centers, it has been proposed that functional disturbances in the vestibular nucleus is an indicator of abnormal activity in the respiratory nucleus<sup>8</sup>.

On the other hand, Gallina et al.<sup>8</sup> reported that the peripheral vestibular system became asymmetric due to hypoxic injury and the peripheral vestibular imbalance was corrected because central vestibular system was more resistant to hypoxic states.

Hypoxic injury in brain stem centers can lead to neurophysiological changes and further deterioration in sleep apnea due to injury in the respiratory centers<sup>4</sup>.

In order to assess brain stem injury in experimental OSAS studies, data from immunohistochemical techniques, magnetic resonance spectroscopy, and audiological tests such as auditory brain stem responses (ABR) have been used<sup>7,26</sup>.

Miceralli et al.<sup>5</sup> showed important impairments in VOR and classical posturographic parameters in patients with OSAS. The mean oxygen saturation has been proposed as a reliable clinical parameter, which indicates progressive disturbance in VOR and vestibulo-spinal reflex.

Mutlu et al.<sup>4</sup> detected abnormalities in brain stem pathways via VEMP recordings in patients with OSAS, and reported that cervical VEMP recordings can be used to detect brain stem damage in severe OSAS.

To our knowledge, no study in the literature has used the BBS to examine fall risk and postural stability in patients with OSAS. In this research, An important limitation was the absence of other vestibular tests (posturography, videonistagmography, VEMP, etc).

In the present study, we detected increased fall risk using BBS in patients with severe OSAS. Further studies investigating postural instability and fall risk in patients with OSAS using the BBS together with other vestibular tests are needed. In conclusion, it can be concluded that the BBS can

contribute in preventing falls (occupational accidents) caused by dizziness in patients with severe OSAS.

### Declaration of interest

Conflicts of interest: none

### Role of the funding source

Source of financial support or funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

### REFERENCES

1. Peppard PE, Young T, Barnet JH, Palta M, Hagen EW, Hla KM. Increased prevalence of sleep-disordered breathing in adults. *Am Journal Epidemiol* 2013;177:1006–1014.
2. Budhiraja R, Parthasarathy S, Quan SF. Endothelial dysfunction in obstructive sleep apnea. *J Clin Sleep Med* 2007;3:409–415.
3. Zhang JH, Fung SJ, Xi M, Sampogna S, Chase MH. Apnea produces neuronal degeneration in the pons and medulla of guinea pigs. *Neurobiol Dis* 2010;40:251–264
4. Mutlu M, Bayır Ö, Yücege MB, Karagöz T, Fırat H, Özdek A, Akın İ, Korkmaz H. Vestibular evoked myogenic potential responses in obstructive sleep apnea syndrome. *Eur Arch Otorhinolaryngol* 2015;272:3137-3141.
5. Micarelli A, Liguori C, Viziano A, Izzi F, Placidi F, Alessandrini M. Integrating postural and vestibular dimensions to depict impairment in moderate-to-severe obstructive sleep apnea syndrome patients. *J Sleep Res* 2017;26:487-494
6. Casale M, Vesperini E, Potena M, Pappacena M, Bressi F, Baptista PJ, Salvinelli F. Is obstructive sleep apnea syndrome a risk factor for auditory pathway? *Sleep Breath* 2012;16:413–417
7. O'Donoghue FJ, Wellard RM, Rochford PD, Dawson A, Barnes M, Ruehland WR, Jackson ML, Howard ME, Pierce RJ, Jackson GD. Magnetic resonance spectroscopy and neurocognitive dysfunction in obstructive sleep apnea before and after CPAP treatment. *Sleep* 2012;35:41–48.
8. Gallina S, Dispenza F, Kulamarva G, Riggio F, Speciale R. Obstructive sleep apnea syndrome (OSAS): effects on the vestibular system. *Acta Otorhinolaryngol Ital* 2010;30:281–284.
9. Yoshida S, Sasa M, Takaori S. Different sensitivity to hypoxia in neuronal activities of lateral vestibular and spinal trigeminal nuclei. *Stroke* 1988;19:357-364.
10. Ferdjallah M, Harris GF, Smith P, Wertsch JJ. Analysis of postural control synergies during quiet standing in healthy children and children with cerebral palsy. *Clin Biomech* 2002;17:203-210.
11. Nashner LM, Black FO, Wall C 3rd. Adaptation to altered support and visual conditions during stance: patients with vestibular deficits. *J Neurosci* 1982;2:536-544.
12. Shumway-Cook A, Horak FB. Assessing the influence of sensory interaction of balance. *Phys Ther* 1986;66:1548-1550.



13. Briggs RC, Gossman MR, Birch R, Drews JE, Shaddeau SA. Balance performance among noninstitutionalized elderly women. *Phys Ther* 1989;69:748–756.
14. Hinman RS, Bennell KL, Metcalf BR, Crossley KM. Balance impairments in individuals with symptomatic knee osteoarthritis: a comparison with matched controls using clinical tests. *Rheumatology* 2002;41:1388-1394.
15. Shumway-Cook A, Brauer S, Woollacott M. Predicting the probability for falls in community-dwelling older adults using the timed up and go test. *Phys Ther* 2000;80:896–903.
16. Baldwin SL, Van Arnam TW, Ploutz-Snyder LL. Reliability of dynamic bilateral postural stability on the Biodex Stability System in older adults, *Med Sci Sport Exerc* 2004;36:530.
17. Aydog ST, Aydog EC, Akci A. Reproducibility of postural stability score in blind athletes. *Isokinetic Exerc Sci* 2004;12:229–232.
18. Gozal D, Kheirandish-Gozal L. Cardiovascular morbidity in obstructive sleep apnea: oxidative stress, inflammation, and much more. *Am J Respir Crit Care Med* 2008;177:369 -375.
19. Fanfulla F, Grassi M, Taurino AE, D'Artavilla Lupo N, Trentin R. The relationship of daytime hypoxemia and nocturnal hypoxia in obstructive sleep apnea syndrome. *Sleep* 2008;31:249-255.
20. Aydog E, Aydog ST, Cakci A. Dynamic postural stability in blind athletes using the biodex stability system. *Int J Sports Med* 2006;27:415-8.
21. Inanır A, Okan S, Filiz B. Evaluation of Postural Stability and Fall Risk in Patients with Rheumatoid Arthritis. *Cukurova Med J* 2013; 38:72-77.
22. Inanır A, Okan S, Filiz B. Evaluation of Postural Stability and Fall Risk in Patients with with Ankylosing Spondylitis. *Cukurova Med J* 2013; 38:86-91.
23. Kayabasi S, Iriz A, Cayonu M, Cengiz B, Acar A, Boynuegri S, Mujdeci B, Eryilmaz A. Vestibular functions were found to be impaired in patients with moderate-to-severe obstructive sleep apnea. *Laryngoscope* 2015;125:1244–1248.
24. Sowerby LJ, Rotenberg B, Brine M, George CF, Parnes LS. Sleep apnea, daytime somnolence, and idiopathic dizziness—a novel association. *Laryngoscope* 2010;120:1274–1278.
25. Quarck G, Ventre J, Etard O, Denise P. Total sleep deprivation can increase vestibulo-ocular responses. *J Sleep Res* 2006;15:369-375.